

**SSI MAXIM**

**TOPSITE™**  
2.5G herbicide

SPECIMEN

**For Use In Noncrop Areas  
And Noncropland Farmsteads**

**ACTIVE INGREDIENTS:**

Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-  
5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid) . . . . . 0.5%  
Diuron (3-[3,4-dichlorophenyl]-1,1-dimethylurea) . . . . . 2.0%

**INERT INGREDIENTS** . . . . . 97.5%  
**TOTAL** . . . . . 100.0%

EPA Reg. No. 34913-22

**KEEP OUT OF REACH OF CHILDREN  
CAUTION! ¡PRECAUCION!**

**PRECAUCION AL USUARIO:** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

See Additional Precautionary Statements  
and Directions For Use Inside



**SSI MAXIM CO., INC.**

P.O. Box 1954 • Kilgore, TX 75663

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

# CAUTION!

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

### FIRST AID

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water.

### PHYSICAL AND CHEMICAL HAZARDS

TOPSITE 2.5G herbicide should be stored only in lined paper bags, stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT store TOPSITE 2.5G herbicide in unlined steel (except stainless steel) containers.

Keep dry.

### ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

### IMPORTANT

DO NOT use on food or feed crops. DO NOT treat irrigation ditches, or water used for crop irrigation or for domestic uses. DO NOT apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result. Keep from contact with fertilizers, insecticides, fungicides and seeds. DO NOT apply or rinse equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts, or similar areas. DO NOT use in California.

### GENERAL INFORMATION

TOPSITE 2.5G herbicide is recommended for control of many annual and perennial grasses and broadleaf weeds on noncropland areas such as railroad, utility, pipeline and highway rights-of-way, petroleum tank farms, pumping installations, fence rows, storage areas, noncropland farmsteads and other similar areas. TOPSITE 2.5G herbicide may be applied at any time throughout the year; however, for maximum effectiveness, the product should be applied prior to weed emergence or when weeds are small. In addition, TOPSITE 2.5G herbicide should be applied prior to ex-

pected rainfall. The degree and duration of control of Kochia and other weed species will depend upon the amount of product applied, the amount of rainfall, soil texture, and other environmental and biological factors.

### DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of SSI MAXIM COMPANY. All such risks shall be assumed by the user.

SSI MAXIM COMPANY warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions of use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values of any other special or indirect damages.

SSI MAXIM COMPANY makes no other express or implied warranty, including other express or implied warranty of FITNESS or MERCHANTABILITY.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** DO NOT contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL FOR BAG:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL FOR PAIL:** Triple rinse (or equivalent). Then offer for recycling or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### APPLICATION INSTRUCTIONS

Uniformly apply TOPSITE 2.5G herbicide with properly calibrated ground equipment, including hand-held spreaders, at a rate of 200 to 300 lbs. product per acre (0.5-0.7 lb/100 sq. ft.).

A preemergent/early postemergent application of TOPSITE 2.5G herbicide will control many weed species, including the following:

## GRASSES

Annual bluegrass (*Poa annua*)  
Bahigrass (*Paspalum notatum*)  
Beardgrass (*Andropogon* spp.)  
Bermudagrass (*Cynodon dactylon*)  
Big bluestem (*Andropogon gerardi*)  
Broadleaf signalgrass (*Brachiaria platyphylla*)  
Browntop millet (*Brachiara ramosa*)  
Canada bluegrass (*Poa compressa*)  
Cattail (*Typha* spp.)  
Centipedegrass (*Eremochloa ophiuroides*)  
Cheat (*Bromus secalinus*)  
Crabgrass (*Digitaria* spp.)  
Dallisgrass (*Paspalum dilatatum*)  
Downy brome (*Bromus tectorum*)  
Fall panicum (*Panicum dichotomiflorum*)  
Fescue (*Festuca* spp.)  
Foxtail (*Setaria* spp.)  
Goosegrass (*Eleusine indica*)  
Guineagrass (*Panicum maximum*)  
Italian ryegrass (*Lolium multiflorum*)  
Johnsongrass (*Sorghum halepense*)  
Kentucky bluegrass (*Poa pratensis*)  
Lovegrass (*Eragrostis* spp.)  
Orchardgrass (*Dactylis glomerata*)  
Paragrass (*Brachiaria mutica*)  
Prairie cordgrass (*Spartina pectinata*)  
Prairie threeawn (*Aristida oligantha*)  
Quackgrass (*Agropyron repens*)  
Sandbur (*Cenchrus* spp.)  
Sand dropseed (*Sporobolus cryptandrus*)  
Smooth brome (*Bromus inermis*)  
Timothy (*Phleum pratense*)  
Torpedograss (*Panicum repens*)  
Vaseygrass (*Paspalum urvillei*)  
Wild barley (*Hordeum* spp.)  
Wild iris (*Iris versicolor*)  
Wild oats (*Avena fatua*)  
Wirestem muhly (*Muhlenbergia frondosa*)  
Witchgrass (*Panicum capillare*)

## BROADLEAF WEEDS

Bigleaf aster (*Aster macrophyllus*)  
Blue mustard (*Chorispora tenella*)  
Bull thistle (*Cirsium vulgare*)  
Burdock (*Arctium* spp.)  
Buttercup (*Ranunculus testiculatus*)  
Camphorweed (*Heterotheca subaxillaris*)  
Canada thistle (*Cirsium arvense*)  
Carpetweed (*Mollugo verticillata*)  
Carolina geranium (*Geranium carolinianum*)

## BROADLEAF WEEDS

Clover (*Trifolium* spp.)  
Cocklebur (*Xanthium strumarium*)  
Common chickweed (*Stellaria media*)  
Common ragweed (*Ambrosia artemisiifolia*)  
Dandelion (*Taraxacum officinale*)  
Dock (*Rumex* spp.)  
Dogfennel (*Eupatorium capillifolium*)  
Eclipta (*Eclipta prostrata*)  
Filaree (*Erodium* spp.)  
Fleabane (*Erigeron* spp.)  
Florida beggarweed (*Desmodium tortuosum*)  
Florida pusley (*Richardia scabra*)  
Giant ragweed (*Ambrosia trifida*)  
Hoary vervain (*Verbena stricta*)  
Horseweed (*Conyza canadensis*)  
Indian mustard (*Brassica juncea*)  
Kochia (*Kochia scoparia*)  
Lambsquarters (*Chenopodium album*)  
Lespedeza (*Lespedeza* spp.)  
Little mallow (*Malva parviflora*)  
Milkweed (*Asclepias* spp.)  
Miners lettuce (*Montia perfoliata*)  
Mullein (*Verbascum* spp.)  
Nettleleaf goosefoot (*Chenopodium murale*)  
Oxeye daisy (*Chrysanthemum leucanthemum*)  
Pepperweed (*Lepidium* spp.)  
Pigweed (*Amaranthus* spp.)  
Plantain (*Plantago* spp.)  
Poison hemlock (*Conium maculatum*)  
Pokeweed (*Phytolacca americana*)  
Primrose (*Oenothera kunthiana*)  
Prostrate knotweed (*Polygonum aviculare*)  
Purslane (*Portulaca* spp.)  
Silverleaf nightshade (*Solanum elaeagnifolium*)  
Smartweed (*Polygonum* spp.)  
Sorrel (*Rumex* spp.)  
Sowthistle (*Sonchus* spp.)  
Sunflower (*Helianthus* spp.)  
Sweet clover (*Melilotus* spp.)  
Tansymustard (*Descurainia pinnata*)  
Texas thistle (*Cirsium texanum*)  
Western ragweed (*Ambrosia psilostachya*)  
White heath aster (*Aster pilosus*)  
Wild carrot (*Daucus carota*)  
Wild lettuce (*Lactuca* spp.)  
Wild parsnip (*Pastinaca sativa*)  
Wild turnip (*Brassica campestris*)  
Woollyleaf bursage (*Ambrosia grayii*)  
Yellow starthistle (*Centaurea solstitialis*)  
Yellow woodsorrel (*Oxalis stricta*)